

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

E83-10012

AgRISTARS

Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

SM-A2-04322

NASA-CR-167648

A Joint Program for
Agriculture and
Resources Inventory
Surveys Through
Aerospace
Remote Sensing

Soil Moisture

May 1982

MEASUREMENT OF SOIL MOISTURE USING REMOTE SENSING MULTISENSOR RADIATION TECHNIQUES

W. P. Waite

(E83-10012) MEASUREMENT OF SOIL MOISTURE
USING REMOTE SENSING MULTISENSOR RADIATION
TECHNIQUES Final Report (Arkansas Univ.)
12 p HC A02/MF A01


N83-12491

CSCI 02C

Unclass
00012

G3/43

Department of Electrical Engineering
University of Arkansas
Fayetteville, Arkansas 72701



Lyndon B. Johnson Space Center
Houston, Texas 77058

1. Report No. SM-A2-04322	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Measurement of Soil Moisture Using Remote Sensing Multisensor Radiation Techniques		5. Report Date May 1982	
		6. Performing Organization Code	
7. Author(s) W. P. Waite		8. Performing Organization Report No.	
		10. Work Unit No.	
9. Performing Organization Name and Address Department of Electrical Engineering University of Arkansas Fayetteville, Arkansas 72701		11. Contract or Grant No. NAS9-14251	
		13. Type of Report and Period Covered Technical	
12. Sponsoring Agency Name and Address AgRISTARS - Soil Moisture Project NASA - Johnson Space Center Houston, Texas 77058		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract The purpose of this study was to investigate means for the microwave remote sensing of soil moisture. The effort was to be directed toward the specification of sensor parameters and the development of inversion algorithms. The conduct of the investigation was a mixture of theoretical modeling, laboratory and field measurement, coupled with analysis of aircraft data obtained from controlled sites.			
17. Key Words (Suggested by Author(s))		18. Distribution Statement	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages	22. Price*

FINAL REPORT

NASA CONTRACT NAS 9-14251

MEASUREMENT OF SOIL MOISTURE USING REMOTE
SENSING MULTISENSOR RADIATION TECHNIQUES

By

W.P. Waite

Department of Electrical Engineering

University of Arkansas

Fayetteville, Arkansas 72701

May 1982

Measurement of Soil Moisture Using Remote Sensing Multisensor Radiation Techniques

INTRODUCTION

The original purpose of this study was to investigate means for the microwave remote sensing of soil moisture. This was both to demonstrate feasibility and to ascertain the effects of perturbing factors such as surface roughness. The effort was to be directed toward the specification of sensor parameters and the development of inversion algorithms. This original goal was subsequently more sharply focussed to concentrate on the development of algorithms specifically useful for agricultural applications, primarily crop yield forecasting.

The conduct of the investigation was a mixture of theoretical modeling, laboratory and field measurement, coupled with analysis of aircraft data obtained from controlled sites. The laboratory and field measurement program was conducted with a bistatic reflectometer. This instrument, while active in nature, was used to measure the specularly reflected component which is more similar to the passive indicimetric measurement than the backscattered component measured by a normal radar. This type operation has the advantage of providing swept (continuous) frequency measurement of the basic radiometric behavior which can only be accomplished using multiple direct frequencies with radiometers.

HISTORICAL SUMMARY

The initial thrust of the program was to evaluate both active (radar) and passive (radiometric) techniques. These were viewed at

ORIGINAL PAGE IS
OF POOR QUALITY

the time not so much as complimentary sensors as competitors. The argument favoring radar centered primarily on its ability to provide fine resolution even at spacecraft altitudes. The principle disadvantage was the known extreme sensitivity to surface roughness. The case for the radiometer was precisely the opposite. It had been observed that the roughness sensitivity was substantially less than that of radar, however, resolution could only be provided by the real antenna beam and would be quite large at spacecraft altitudes.

The initial investigations at the University of Arkansas concentrated on the effects of surface roughness as a function of frequency and the analysis of passive data available from the first aircraft missions. The investigation of roughness concentrated on defining the variation of the specularly reflected (coherent) component as a function of frequency (Hancock, 1977). The hope was to use multiple (continuous) frequency measurement to define the rate of decay with frequency of the specularly reflected component. Once this rate was defined, it would be possible to invert to an equivalent smooth surface reflectivity from which the determination of dielectric constant and then soil moisture would be a relatively simple matter.

It should again be noted that although the measurement conducted utilized active swept frequency techniques the implementation method would be multiple frequency passive since it is desired to monitor the coherent component of the scattered signal. This is possible, of course, only over the range where the incoherent (diffuse) component may be neglected in comparison to the coherent component. This study and continuing investigations of rough surface effects generated the following conclusions:

ORIGINAL PAGE 13
OF POOR QUALITY

- 1) Correction for roughness can be accomplished, however, even for frequencies as low as 1GHz , the effective value of the rms height that can be accommodated is significantly less than $1/\text{cm}$.
- 2) The effective roughness of real surfaces with discontinuities is substantially greater than their measured rms height.
- 3) The measurements exhibited a marked sensitivity to changes in the correlation distance of the surface.

These observations could be further summarized to say that multi-parameter measurement of the exponential decrease in the coherent scattered component does not appear feasible for realistic agricultural surfaces unless frequencies substantially below 1GHz are used.

These conclusions still have significance today in that they imply that the observed decreased sensitivity due to roughness of the passive versus active sensors is due to the fact that the passive system is responding to the diffuse component of the scattered signal. This in turn implies that the exponential correction factor applied to explain the sensitivity compression due to roughness is incorrect. The correction should be a reduction from the plane surface value to the diffuse value since for the measured data the diffuse component is significantly greater than the specular component.

The analysis of airborne passive radiometric data presented an entirely different problem. The difficulty here was not in showing a pronounced sensitivity due to soil moisture, but in interpreting the significance of the results. The only means of evaluating the data is by linear correlation analysis. The results consistently

ORIGINAL PAGE 13
OF POOR QUALITY

demonstrated a slightly higher correlation with the upper levels of the moisture profile both at X and L band. However, the internal correlation of the profile also leads to correlation of brightness temperature with moisture content at depths well below any reasonable expectation of penetration.

At approximately this time the emphasis of the program was sharply focussed on the application to crop yield estimation. This immediately led to consideration of means to estimate the total plant water availability which in turn leads to requiring a knowledge of the profile to depths determined by root density which may range to six feet or more. Thus, while the measurement technique demonstrated increasingly good correlation with approximately the upper one inch of the surface the application suddenly required estimation to six feet.

A study of soil moisture budget models was conducted to assess the feasibility of periodic surface measurements to update a moisture budget model driven by meteorologic data. This, in essence, periodically corrects the budget model by enforcing a measured boundary condition.

Considerations such as this led to the need for maximizing the actual measurement depth. While there is no hope of performing measurements to the depth required by the application, the greater the depth that can be sampled, the greater will be the accuracy and the longer can be the period between measurements.

At approximately this time it could be concluded that microwave remote measurement of the soil moisture of bare surfaces was certainly feasible. In fact, it appeared this could be done quite accurately

**ORIGINAL PAGE IS
OF POOR QUALITY**

even with the perturbations introduced by surface roughness. The problem lie in determining the depth to which the moisture might be sensed and defining at least a moisture gradient for the upper surface. In line with this a program of laboratory measurements of artificially constructed and layered moisture profiles was conducted. This program definitely has established the ability to detect discontinuities in moisture to significant depths (greater than six inches). The problem with such artifical profiles was that although the layer interference pattern could be used to accurately measure depth and infer moisture content, it likewise required continuous frequency measurement.

The next step in the process was to conduct a series of field measurements over controlled moisture plots to monitor the effects of natural gradients under controlled conditions. This program offered as well the first opportunity to compare the microwave measurements with soil matric potential rather than with soil moisture. As had been predicted this offered improved correlation.

The field measurement program demonstrated that naturally occurring moisture gradients in the upper surface could in fact yield the effect of a subsurface layer. While for each individual frequency over the range the reflectivity tracks the decrease in moisture with evaporation, the continuous frequency trace may exhibit an interference pattern. The coherent component transmitted into the surface and leading to the interference effect is apparently decreased by the surface roughness. Thus, even natural surfaces, unless sufficiently rough, will produce pronounced coherent effects introducing a scatter into the measurement.

Also near the end of this program data became available from the Colby test site. The initial analysis of these data confirmed the measurements of SCZ showing row orientation to have a pronounced effect upon the active backscatter return even at K Land. This was in distinct contradiction to the conclusions reached from the measurements of the University of Kansas MAS system from which it was concluded that row effects would be insignificant at C band.

CONCLUSIONS

The preceding has been a brief synopsis of the studies conducted at the University of Arkansas. These were, of course, simply part of an overall program including primarily NASA/JSC, NASA Goddard, the University of Kansas, and Texas A & M University as well. This program has undoubtedly aided substantially in the understanding of the microwave response due to soil moisture. As one example of this the development of models to predict the complex dielectric constant have been produced and have lead to interpretation of the results in terms of a matric potential rather than simply moisture content. This appears to have both a better theoretical basis and likewise to be the parameter of more concern to the user as it is directly related to the water available to the plant. Similar advances have been made in the development of coherent and incoherent radiative transfer models and rough surface scattering models.

While these advances in understanding have been substantial they have not as yet answered the question of whether satellite remote sensing of soil moisture is feasible or practical for agricultural and hydrologic uses. The greatest single problem is not lack of demonstrated sensitivity but definition as to what moisture the

ORIGINAL PAGE IS
OF POOR QUALITY

instrument is responding. With the internal correlation of virtually any moisture profile, a limited data sample may demonstrate excellent correlation at even substantial depths. It still remains to determine the best inversion algorithm for moisture input to budget models over a wide range of profile shapes.

The degree of the row orientation effect is still to be completely specified and may well force active systems to adopt a cross polarized measurement. The effect of screening vegetation remains somewhat controversial with a wide range of results from differing measurement programs. There is still some disagreement as to the severity of coherent effects for differing profile shapes particularly in the near surface. Efforts in this area are hampered by a lack of data for near surface profiles particularly with the depth resolution needed.

In final conclusion it would appear that the advances in the understanding of microwave soil moisture measurement have been substantial. These have been for the most part accomplished with relatively isolated data sets from both ground measurement and aircraft programs. It would appear that perhaps the most pressing need is for measurement programs relatively small in area but providing continuous measurement over an extended period of time. This should not be surprising as agriculturists have used this approach for years due to the great variability encountered. Since the soil is the object studied here the same method would seem to apply.

ORIGINAL PAGE IS
OF POOR QUALITY

REPORTS AND PUBLICATIONS

The basic reporting mechanism during the conduct of this program was presentation of results at quarterly review sessions held at JSC. Approximately 20 of these presentations were made with the delivery of all presentation material and descriptive synopsis. These formed the basis for directing the program and most importantly for rapid interaction between all members of the investigative team. Publications and in depth reports on specific aspects are listed below:

Waite, W.P., "Frequency Dependence of the Microwave Brightness
Temperature of Rough Dielectric Surfaces,"
Proc. Am. Soc. of Photogrammetry, Little Rock, AR, pp. 377-392,
ct. 1977.

Waite, W.P., et al., "Long Wavelength Radar for Geological Analysis in
Vegetated Terrain," Proc. Am. Soc. of Photogrammetry,
Albuquerque, NM, pp. 386-39A, Oct. 1978.

Waite, et al. "Dual Polarized Long Wavelength Radar for Discrimination
of Agricultural Land Use," Proc. Am. Soc. of Photogrammetry,
Albuquerque, NM, pp. 595-607.

Waite, W. "Aircraft Experiments Required for Development of Microwave
Soil Moisture Measurement," NASA Hdqtrs. Joint Soil Moisture
Committee Meeting, Washington, DC, Jan. 1979.

Waite, W.P. et al. "Data Documentation for the Bare Soil Experiment at
the University of Arkansas." p. 127, Jan. 1980.

Waite, W.P. et al., "Use of Satellite Radar Imagery for the Detection
of Standing Water Beneath Forest Vegetation,"
Proc. Amer. Soc. Photogramm., Niagara Falls, NY, Oct. 1980.

ORIGINAL PAGE IS
OF POOR QUALITY

- Waite, W.P. et al., "Look Direction Dependence of Radar Backscattering Cross Section for Agricultural Fields,"
Proc. Amer. Soc. Photogramm., Niagara Falls, NY, Oct. 1950.
- Stiles, W. "The Effect of Surface Roughness on the Microwave Measurement of Soil Moisture Content," MS Thesis, University of Arkansas, 1974.
- Turner, C.H., "The Effect of Low Level Vegetation on the Microwave Measurement of Soil Moisture Content," MS Thesis, University of Arkansas, 1974.
- Moss, R.A., "Analysis of Airborne Passive Microwave Data for the Remote Measurement of Soil Moisture," MS Thesis, University of Arkansas, 1977.
- McCall, E./M., "Analysis of Radiative Transfer Model for the Prediction of Microwave Emission and Soil Moisture," MS Thesis University of Arkansas, 1977.
- Hancock, G.D., "Broad Spectrum Microwave Measurement of Rough Discontinuous Surfaces for the Determination of Moisture Content," MS Thesis, University of Arkansas, 1977.
- Corball, B.H., "The Effects of Row Orientation and Soil Moisture on Radar Backscattering from Agricultural Fields," MS Thesis, University of Arkansas, 1980.
- Reid, C.S., "The Utility of Microwave Remote Sensing Techniques for Soil Moisture Profile Determination," MS Thesis, University of Arkansas, 1977.
- Hancock, G.D., "Inversion Algorithms for the Microwave Remote Sensing of Soil Moisture, Ph.D. Dissertation, University of Arkansas, 1982.